

**SDC Slovenia**  
Clinical Development



# **BCS3 compounds: In Vivo experience with non-Q1/Q2 formulations**

**Igor Legen, Ph.D.**

**Head Clinical Development**

**Sandoz Development Center Slovenia, Lek Pharmaceuticals d.d.**

**SANDOZ** A Novartis  
Division

2021 Generic Drug Science and Research Initiatives Public Workshop, June 23, 2021

# Disclaimer

„The opinions expressed herein are solely those of the presenter and do not represent statements or opinions of Lek Pharmaceuticals d.d., Sandoz Pharmaceuticals d.d. or Novartis Pharma Services Inc.« “

# Background

## Biopharmaceutics Classification System (BCS) <sup>1</sup>

- Class I: high solubility, high permeability
- Class II: low solubility, high permeability
- Class III: high solubility, low permeability
- Class IV: low solubility, low permeability

# Background

In vivo bioequivalence studies may be waived for immediate release formulations with not narrow therapeutic index BCS1 and BCS3 compounds provided <sup>2</sup>:

- Dissolution: BCS1 - rapid or BCS3 - very rapid
- Adequate stability in GIT
- Excipients similarity:
  - BCS1: excipients that may affect absorption - qualitatively the same and quantitatively similar
  - BCS3: all excipients qualitatively the same and quantitatively very similar

# Excipients

Impact on :

- permeability across intestinal epithelium <sup>3</sup>:
  - tight junctions
  - membrane fluidity
  - transporters
- intestinal transit<sup>10</sup>

Permeability: Caco-2 cells - more sensitive compared to native intestinal tissue and to In Vivo results <sup>4,5</sup>.

Small quantities of certain excipients can influence bioavailability of some compounds <sup>6</sup>.

# Bioequivalence risk

Are bioequivalence studies with BCS3 compounds more risky ?

Source	Success of bioequivalence studies (%)		
	BCS1	BCS3	BCS2
Lamouche S et al., <sup>7</sup>	89	89	72
Cristofolletti R et al., <sup>8</sup>	84	90	60
Ramirez E et al., <sup>9</sup>	85	85	50


# Sandoz experience with BCS3 studies (10+ years)

Bioequivalence studies with immediate release BCS3 formulations:

- 16 INNs
- 25 different formulations
- 83 studies
  - 34 with mono and 49 with combo product
  - 66 under fasting and 17 under fed condition
  - 70 (84 %) qualitatively **different** formulation as reference (non-functional coating excluded)
  - 82 (99 %) successful studies

# Case 1: Presence of „critical“ excipient

- Immediate release formulation, BCS3 compound
  - Reference: Compound
  - Test: Compound in the form of Hydroxypropyl- $\beta$ -cyclodextrin complex (HP-  $\beta$ -CD)



Viability:

- Isc
- TEER
- PD

Rat jejunum, Side-by-side diffusion chamber, pH=6.85, 37°C

	Compound	Compound - HP- $\beta$ -CD complex
$P_{APP}$ ( $\times 10^{-6}$ cm/s)	$3.38 \pm 0.22$	$3.51 \pm 0.18$
TEER ( $\text{ohm} \times \text{cm}^2$ )	$32.9 \pm 5.4$	$34.5 \pm 2.5$
PD (mV)	$-1.11 \pm 0.13$	$-1.03 \pm 0.09$
Isc ( $\mu\text{A}/\text{cm}^2$ )	$34.7 \pm 2.7$	$31.2 \pm 4.4$

N=3, average  $\pm$  SEM



# Case 1: Presence of „critical“ excipient

Comparative, Single-Dose, 2-way Crossover Bioavailability Study in 36 Healthy Adult Male Volunteers under Fasting Conditions


Comparison	Parameter	Ratio	Lower 90% CI	Upper 90% CI
T vs R	C <sub>max</sub>	94	88	100
T vs R	AUC <sub>t</sub>	102	98	106
T vs R	AUC <sub>i</sub>	102	99	106



# Case 2: Q2 difference in „critical“ excipient

- Granules for oral solution, BCS1/2 compound (salt of an acidic drug)
  - Reference: 1.70 g mannitol
  - Test: 1.82 g mannitol

7% difference



2% difference in SIT

## Case 2: Q2 difference in „critical“ excipient

Comparative, Single-Dose, 2-way Crossover Bioavailability Study in 26 Healthy Adult Volunteers under Fed Conditions

Comparison Parameter		Ratio	Lower 90% CI	Upper 90% CI	
T vs R	Cmax	105	97	113	✓
T vs R	AUCt	96	94	99	✓

# Conclusions

- BCS1 and BCS3 compounds bring similar bioequivalence risk to IR formulations
- Q1 difference in excipients does not represent a major risk for bioequivalence for BCS3 containing IR formulations
- Impact of Q1 and Q2 difference in „critical“ excipients on drug absorption should be evaluated on a case by case basis

# References

- <sup>1</sup> Amidon GL et al., 1995, A Theoretical Basis For a Biopharmaceutics Drug Classification: The Correlation of In Vitro Drug Product Dissolution and In Vivo Bioavailability, *Pharm Res*, 12: 413-420.
- <sup>2</sup> ICH M9 Guideline on biopharmaceutics classification system-based biowaivers, ICH, (adopted by EMA 2020 and by FDA 2021)
- <sup>3</sup> Aungst JB, 2000, Intestinal Permeation Enhancers, *J Pharm Sci*, 89: 429-446.
- <sup>4</sup> Legen I et al., 2005, Comparison of different intestinal epithelia as models for absorption enhancement studies, *Int J Pharm*, 291: 183-188.
- <sup>5</sup> Vaithianathan S et al., 2016, Effect of Common Excipients on the Oral Drug Absorption of Biopharmaceutics Classification System Class 3 Drugs Cimetidine and Acyclovir, *J Pharm Sci*, 105: 996-1005.
- <sup>6</sup> Garcia-Arieta A, 2014, Interactions between active pharmaceutical ingredients and excipients affecting bioavailability: Impact on bioequivalence, *Eur J Pharm Sci*, 65: 89-97.
- <sup>7</sup> Lamouche S et al., 2008. The biopharmaceutical classification system: Can it help predict bioequivalence outcome? A CRO retrospective analysis, *AAPS\_abstract* 002992
- <sup>8</sup> Cristofolletti R et al., 2013, A Comparative Analysis of Biopharmaceutics Classification System and Biopharmaceutics Drug Disposition Classification System: A Cross-Sectional Survey with 500 Bioequivalence Studies, *J Pharm Sci*, 102: 3136-44.
- <sup>9</sup> Ramirez E et al., 2010, Acceptability and characteristics of 124 human bioequivalence studies with active substances classified according to the Biopharmaceutic Classification System, *Br J Clin Pharmacol*, 70: 694-702.
- <sup>10</sup> Adkin DA et al., 1995, The effect of different concentrations of mannitol in solution on small intestinal transit: Implications for drug absorption, *Pharm Res*, 12: 393-396.

# Acknowledgment

Sandoz Product Development

- Clinical Development
- IVIVC



**Thank you**